

First Genetic Data of *Nebalia koreana* (Malacostraca, Leptostraca) with DNA Barcode Divergence among *Nebalia* Species

Ji-Hun Song^{1,2}, Gi-Sik Min^{1,*}

¹Department of Biological Sciences, Inha University, Incheon 22212, Korea

²Animal & Plant Resources Research Division, Nakdonggang National Institute of Biological Resources,
Sangju 37242, Korea

ABSTRACT

We determined the cytochrome *c* oxidase subunit 1 (*COI*) sequences of *Nebalia koreana* Song, Moreira & Min, 2012 (Leptostraca) collected from five locations in South Korea, and this represents the first genetic data of this species. The maximum intra-species variation was 1.2% within *Nebalia hessleri* Martin, Vetter & Cash-Clark, 1996, while inter-species variation ranged from 9.0% (*N. hessleri* and *Nebalia gerkenae* Haney & Martin, 2000) to 34.8% (*N. hessleri* and *Nebalia pseudotronicosoi* Song, Moreira & Min, 2013). This result is well agreed with the interspecific relationships among *Nebalia* species based on morphological characteristics. In conclusion, this study showed the usefulness of *COI* sequences as a DNA barcode within the genus *Nebalia* Leach, 1814.

Keywords: *COI*, DNA barcode, Leptostraca, *Nebalia*, South Korea

INTRODUCTION

The order Leptostraca Claus, 1880 is the only extant order in the subclass Phyllocarida Packard, 1879, and is considered by many researchers to be the most primitive group in the class Malacostraca Latreille, 1802 (Claus, 1888; Manton, 1934; Hessler and Newman, 1975; Barnes, 1980; Dahl, 1987; Walker-Smith and Poore, 2001). The genus *Nebalia* Leach, 1814 is the largest member of the order Leptostraca (38 of 60 valid species), and so far, the following three species have been reported in South Korea: *Nebalia dolsandoensis* Song & Min, 2017, *Nebalia koreana* Song, Moreira & Min, 2012, and *Nebalia pseudotronicosoi* Song, Moreira & Min, 2013 (see Song et al., 2012, 2013; Song and Min, 2017). *Nebalia koreana* is distributed in Eastern, Western and Southern coastal waters of Korea, while the other two species were found restrictively at their type localities (Song, 2013). *Nebalia koreana* is diagnosed by the following characteristics: antennule fourth article with up to five short distal spines; pleonites 3–7 with rounded denticles on posterior dorsal margins; and pleopod 4 protopod lacking serrations on posterior margin.

Among the mitochondrial genes already examined in most animal phyla, including Crustacea, the cytochrome *c* oxidase subunit 1 (*COI*) gene has proved to be a particularly useful taxonomic marker (Meyran et al., 1997; Wares, 2001; Hajibaei et al., 2006; Clare et al., 2007; Elsasser et al., 2009; Zemlak et al., 2009). However, the usefulness of the *COI* sequences in *Nebalia* species has never been proven. In this study, we firstly determined the *COI* sequence of *N. koreana* (Fig. 1), and attempted to verify its usefulness.

RESULTS AND DISCUSSION

The *COI* sequences of *N. koreana* were obtained from specimens collected from each of the following five locations in South Korea: Bogildo (34°08'10"N, 126°33'48"E), Cheongsando (34°10'54"N, 126°33'59"E), Mokpo (34°48' 19"N, 126°21'43"E), Masan (35°11'11"N, 128°33'53"E), and Sokcho (38°11'50"N, 128°35'00"E) using light traps according to the methods described by Song et al. (2013). The morphological identification, genomic DNA extraction, and PCR conditions were performed according to the methods de-

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

***To whom correspondence should be addressed**

Tel: 82-32-860-7692, Fax: 82-32-874-6737
E-mail: mingisik@inha.ac.kr



Fig. 1. *Nebalia koreana* collected from Sokcho, lateral view.

Table 1. The minimum/maximum values of inter-species pairwise variation (%) between *COI* sequences from *Nebalia* species

	1	2	3	4	5
1. <i>N. koreana</i>	–				
2. <i>N. dolsandoensis</i>	16.4/16.7	–			
3. <i>N. pseudotronicosoi</i>	16.8/17.1	18.7/18.7	–		
4. <i>N. gerkenae</i>	30.8/31.5	29.7/30.4	32.5/32.5	–	
5. <i>N. hessleri</i>	32.2/34.0	31.1/31.9	34.4/34.8	9.0/10.3	–

The GenBank accession numbers used in this study are as follows. *Nebalia koreana*, MK396103–MK396107 (this study); *N. dolsandoensis*, KT229637 and KT229638; *N. pseudotronicosoi*, JX442539–JX442543; *N. gerkenae*, FJ170145–FJ170147; *N. hessleri*, FJ170162–FJ170169

scribed by Song and Min (2017). The sequences were aligned using Geneious 8.1.9 (Biomatters Ltd., Auckland, New Zealand). The *COI* sequences determined from this study and retrieved from GenBank were aligned using MEGA 10 (Kumar et al. 2018). Nucleotide sequence divergence was calculated using the Kimura two-parameter distances.

In our analysis, the maximum intra-species variation was 1.2% within *Nebalia hessleri* Martin, Vetter & Cash-Clark, 1996, while inter-species variation ranged from 9.0% (*N. hessleri* and *Nebalia gerkenae* Haney & Martin, 2000) to 34.8% (*N. hessleri* and *N. pseudotronicosoi*) (Table 1). This result is well agreed with the interspecific relationships among *Nebalia* species based on morphological characteristics (Song and Min, 2017). In conclusion, this study suggests that the *COI* sequences are very useful for the identification of *Nebalia* species, as confirmed by many other crustaceans.

ACKNOWLEDGMENTS

This work was supported by grant from Inha University.

REFERENCES

- Barnes RD, 1980. Invertebrate zoology. Saunders College Publishing, Philadelphia, pp. 1-1089.
- Clare EL, Lim BK, Engstrom MD, Eger JL, Hebert PDN, 2007. DNA barcoding of Neotropical bats: species identification and discovery within Guyana. *Molecular Ecology Notes*, 7:184-190. <https://doi.org/10.1111/j.1471-8286.2006.01657.x>
- Claus C, 1888. Über den Organismus der Nebaliden und die systematische Stellung der Leptostraken. *Arbeiten aus dem Zoologischen der Universität*, 8:1-148.
- Dahl E, 1987. Malacostraca maltreated: the case of the Phyllocarida. *Journal of Crustacean Biology*, 7:721-726. <https://doi.org/10.1006/jcrub.1987.0011>

- doi.org/10.1163/193724087X00469
- Elsasser SC, Floyd R, Hebert PDN, Schulte-Hostedde AI, 2009. Species identification of North American guinea worms (Nematoda: Dracunculus) with DNA barcoding. *Molecular Ecology Resources*, 9:707-712. <https://doi.org/10.1111/j.1755-0998.2008.02393.x>
- Hajibaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN, 2006. DNA barcodes distinguish species of tropical Lepidoptera. *Proceedings of the National Academy of Sciences of The United States Of America*, 103:968-971. <https://doi.org/10.1073/pnas.0510466103>
- Hessler RR, Newman WA, 1975. A trilobitormorph origin for the Crustacea. *Fossils and Strata*, 4:437-459.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K, 2018. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution*, 35:1547-1549. <https://doi.org/10.1093/molbev/msy096>
- Manton SM, 1934. On the embryology of the crustacean *Nebalia bipes*. *Philosophical Transactions of the Royal Society of London*, 223:163-238. <https://doi.org/10.1098/rstb.1934.0005>
- Meyran JC, Monnerot M, Taberlet P, 1997. Taxonomic status and phylogenetic relationships of some species of the genus *Gammarus* (Crustacea, Amphipoda) deduced from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution*, 8:1-10. <https://doi.org/10.1006/mpev.1996.0399>
- Song JH, 2013. A systematic study on the order Leptostraca (Crustacea: Malacostraca) from Korea. MS thesis, Inha University, Incheon, Korea, pp. 1-96.
- Song JH, Min GS, 2017. A new species of *Nebalia* (Malacostraca: Phyllocarida: Leptostraca) from South Korea, with a key to the species of *Nebalia* Leach, 1814. *Journal of the Marine Biological Association of the United Kingdom*, 97:59-68. <https://doi.org/10.1017/S0025315415002180>
- Song JH, Moreira J, Min GS, 2012. A new species of Leptostraca, *Nebalia koreana* (Malacostraca: Phyllocarida), from South Korea. *Journal of Crustacean Biology*, 32:641-653. <https://doi.org/10.1163/193724012X638482>
- Song JH, Moreira J, Min GS, 2013. *Nebalia pseudotronicosoi* n. sp. (Malacostraca: Leptostraca), from South Korea, with a peculiar sexual dimorphism. *Journal of Crustacean Biology*, 33:124-136. <https://doi.org/10.1163/1937240X-00002106>
- Walker-Smith GK, Poore GCB, 2001. A phylogeny of the Leptostraca (Crustacea) with keys to families and genera. *Memoirs of Museum Victoria*, 58:383-410. <https://doi.org/10.24199/j.mmv.2001.58.21>
- Wares JP, 2001. Patterns of speciation inferred from mitochondrial DNA in North American *Chthamalus* (Cirripedia: Balanomorpha: Chthamaloidea). *Molecular Phylogenetics and Evolution*, 18:104-116. <https://doi.org/10.1006/mpev.2000.0857>
- Zemlak TS, Ward RD, Connell AD, Holmes BH, Hebert PDN, 2009. DNA barcoding reveals overlooked marine fishes. *Molecular Ecology Resources*, 9:237-242. <https://doi.org/10.1111/j.1755-0998.2009.02649.x>

Received January 9, 2019
 Revised January 14, 2019
 Accepted January 15, 2019